

The role of perinatal problems in risk of co-morbid psychiatric and medical disorders in adulthood

YOKO NOMURA^{1*}, JEANNE BROOKS-GUNN^{2,3}, CHARLES DAVEY⁴, JACOB HAM⁵
AND WILLIAM P. FIFER^{3,6,7}

¹ Mount Sinai School of Medicine, New York, NY, USA; ² Teachers College, Columbia University, New York, USA; ³ Columbia University College of Physicians and Surgeons, New York, USA; ⁴ The School at Columbia University, New York, USA; ⁵ Beth Israel Medical Center and Albert Einstein College of Medicine, New York, NY, USA; ⁶ New York State Psychiatric Institute, New York, NY, USA; ⁷ Sackler Institute for Developmental Psychobiology, New York, NY, USA

ABSTRACT

Background. Perinatal problems may be associated with an increased risk for psychological and physical health problems in adulthood, although it is unclear which perinatal problems (low birthweight, preterm birth, low Apgar scores, and small head circumference), or what clusters of problems, are more likely to be associated with later health problems. It is also not known whether perinatal problems (singly or together) are associated with co-morbidity between psychological and physical health problems.

Method. A regional random sample (from Baltimore) of mothers and their children ($n = 1525$) was followed from birth to adulthood (mean age 29 years). Perinatal conditions were measured at delivery. Psychological problems (depression and suicidal ideation) were measured with the General Health Questionnaire-28 (GHQ-28) and physical problems (asthma and hypertension) with the RAND-36 Health Status Inventory.

Results. Children with perinatal problems were generally at increased risk for depression, suicidal ideation and hypertension, and co-morbid depression and hypertension even after controlling for confounders. One possible underlying condition, preterm low birthweight (LBW), extracted by cluster analysis, considering all of the four perinatal problems, was associated with increased risk for psychological and physical health outcomes as well as co-morbidity of the two.

Conclusions. LBW, preterm birth and small head circumference singly increased the risk for both psychological and physical health problems, as well as co-morbid depression and hypertension, while low Apgar scores were only associated with psychological problems. Delineating different etiological processes, such as preterm LBW, considering various perinatal problems simultaneously, might be of benefit to understanding the fetal origin of adult illness and co-morbidity.

INTRODUCTION

The high rate of psychological and physical health problems is a major public health issue. People with co-morbid depression and physical health problems have increased functional impairment (Lee *et al.* 2000), absenteeism

(Broadhead *et al.* 1990), hospital stays (Fulop *et al.* 1988), and suicidal ideation (Druss & Pincus, 2000). The physical and psychological suffering of individuals with co-morbid depression and other physical health problems is synergistically greater than that of either problem alone.

There is mounting evidence that a higher prevalence of co-morbid psychological problems, such as depression, exists among persons with heart disease (Anda *et al.* 1993; Musselman

* Address for correspondence: Yoko Nomura, Ph.D., M.P.H., Department of Psychiatry, Mount Sinai School of Medicine, One Gustav L. Levy Place, Box 1230, New York, NY 10029, USA.
(Email: yoko.nomura@mssm.edu)

et al. 1998; Cox *et al.* 2002; van Melle *et al.* 2004; Glassman *et al.* 2006) and respiratory problems (Wells *et al.* 1988; Hurwiz & Morgenstern, 1999; Vila *et al.* 2000; Wamboldt *et al.* 2000; Gillaspay *et al.* 2002; Goodwin *et al.* 2004; Ortega *et al.* 2004). However, it is still unclear whether the co-morbidity is caused by one or other illness, or whether even a third factor might underlie the association (Kramer *et al.* 1998).

One such third factor might be perinatal risk, such as low birthweight (LBW) and prematurity (Allen *et al.* 1998; Wadhwa *et al.* 2001). A review of studies on perinatal risk and later health problems suggested that exposure to stress *in utero* and/or early in life may increase vulnerability to subsequent normative stress and impair functions of the central nervous system (CNS) and hypothalamic–pituitary–adrenal (HPA) axis (Sanchez *et al.* 2001). These changes are hypothesized to elevate the risk for both emotion regulation and physical health problems, particularly related to respiratory function (Wamboldt *et al.* 2000) and cardiovascular reactivity (Sanchez *et al.* 2001). Animal studies show that chronic perinatal stress increases the risk for subsequent problems in physical health, affect and behavior (Ward, 1991). Several aversive experiences *in utero* and in the neonatal period result in a sensitization of emotional and HPA-axis responses to subsequent stress (Chuckley, 1996; Meaney, 2001). Human studies similarly suggest that stressful prenatal experiences, such as maternal smoking, preterm birth or LBW, may contribute to abnormalities in the endocrine system (Wadhwa *et al.* 1998; Heim & Nemeroff, 2001). This pathway may be mediated by increased exposure to corticotropin-releasing hormone (CRH), because CRH is involved in the organization of the CNS during this sensitive period (Sandman *et al.* 1998).

Despite its potential importance, this third risk factor remains largely unexplored. Studies examining psychological and physical health problems are generally conducted separately and not integrated into a broader, potentially more constructive, perspective. Furthermore, many studies of perinatal risk factors measure markers of perinatal problem, such as LBW, preterm birth, small head size and low Apgar scores, but their underlying causes and their long-term consequences could be different. For

example, LBW primarily caused by preterm birth has been associated with an elevated risk for cardiovascular illness (Leon *et al.* 1998; Barker *et al.* 2002; Huxley *et al.* 2002; Davies *et al.* 2006), while being born small for gestational age (SGA) has been associated with an increased risk for diabetes (Hales *et al.* 1991; Veening *et al.* 2002; Vaag *et al.* 2006).

To test this third factor hypothesis, we have analyzed data from the Johns Hopkins Collaborative Perinatal Study (JHCPS). Using the same data, Fan & Eaton (2001) reported a three- to fourfold increased risk for general mental distress and nervous conditions in adulthood among those with perinatal risk, such as LBW and preterm birth, and this risk was accentuated in the low income group. However, they did not examine whether the same perinatal problems increased the risk for physical health problems alone or in conjunction with psychological problems. It would be fruitful if we could not only estimate the risk for adult health problems by perinatal indicators singly but also distinguish different underlying etiological processes, taking advantage of the available perinatal indicators (LBW, preterm birth, small head circumference and low Apgar scores).

In the study reported here, we examined whether four perinatal risk indicators singly increase the risk for psychological problems (e.g. depression; Hulshoff *et al.* 2000; Thompson *et al.* 2001; Patton *et al.* 2004; Nomura *et al.* in press) and physical health problems (e.g. respiratory and cardiovascular problems; Barker & Lackland, 2003; Eriksson *et al.* 2001). We also examined the magnitude of the co-morbidity between the two. Finally, we attempted to elucidate the different underlying perinatal etiological processes and their effects on adult health. Given our underlying assumption that general perinatal problems may lead to biophysiological changes, associated with both psychological problems (related to emotional regulation) and physical health problems (related to cardiovascular and respiratory functions), we focused primarily on depression and suicidal ideation as our psychological outcomes and hypertension and asthma as our physical health outcomes. We hypothesized that (1) perinatal problems are associated with an increased risk for psychological and physical health problems; (2) there would be an elevated

risk for depression or suicidal ideation if hypertension or asthma was also present; (3) perinatal problems are associated with an increased risk for co-morbid psychological and physical problems; and (4) different underlying etiological processes may exist and the risks may differ depending on the process.

METHOD

Data source and participants

The JHCPS, part of the National Collaborative Perinatal Project (NCPP), consists of prospective data collected from a random sample of pregnant women who received prenatal care, and delivered their babies in hospital during 1960–1964. The NCPP entailed a single study design across all sites and involved the systematic data collection from pregnancy to the first 8 years to identify perinatal and early childhood factors that adversely affect subsequent child development (Fan & Eaton, 2001). Reports from the NCPP have been summarized elsewhere (Nichols & Chen, 1981; Hardy, 2003).

In 1992–1994, the Pathways to Adulthood Study at Johns Hopkins University (principal investigators: J. Hardy & S. Shapiro) bridged the period from age 7–8 to 27–33 years of age. Of the 2694 offspring eligible for the Pathways to Adulthood Study, 2220 (82%) were located and 1758 completed interviews (71.4% response rate of known outcome and 65.3% of full interview). Those who were located but did not complete full interviews had mothers with characteristics generally similar to those who were interviewed. The study design, interview procedure, potential biases and attrition of the sample are described in detail elsewhere (Hardy *et al.* 1997). Of the 1758 offspring who completed interviews, 1525 provided information on perinatal factors and psychological and physical health problems in adulthood. This subset provided the data used in this paper. Analyses of demographic and perinatal differences between the 1525 included and 1069 excluded offspring revealed no difference in race (African American: 81.9% *v.* 81.3%), birthweight (3036 *v.* 3009 g), gestational age (38.9 *v.* 38.6 weeks), head circumference (33.54 *v.* 33.46 cm) and 5-min Apgar scores (8.83 *v.* 8.79). However, the included offspring were more likely to be female than the excluded (54.4% *v.* 43.4%, $p < 0.001$).

This study was ruled exempt by the Institutional Review Board Committee at Mount Sinai School of Medicine because it involved secondary data analysis of de-identified data.

Measures

Perinatal problems

Independent variables included four perinatal problems. Birthweight, 5-min Apgar scores and head circumference were recorded by a research nurse in the delivery room. Gestational age was determined by a combination of the mother's self-report of their last menstrual period and sonogram. We used both continuous and dichotomous measures of perinatal indicators. For dichotomous variables, we used the traditional cut-off points to create risk indicators for LBW (<2500 g), low 5-min Apgar score (<7), small head circumference (<32 cm), and preterm birth (<37 weeks' gestational age).

Adult psychopathology and illness

Depression was measured with the General Health Questionnaire-28 (GHQ-28; Goldberg, 1978). It was assessed by responses to seven questions, with response options ranging from 1 (better than usual) to 4 (much worse). Using the scoring method, a choice of 1 or 2 was recoded as '0' and 3 or 4 as '1'. Four or more '1' responses were used to indicate the presence of depression. The internal consistency of the GHQ is high (Goldberg, 1972). Compared to the three most commonly used instruments for identifying psychiatric illness (the Center for Epidemiological Studies Depression Scale, the Beck Depression Inventory, and the Hospital Anxiety and Depression Scale), the GHQ had higher sensitivity (92%) and specificity (90%) for identifying psychiatric illness (Clarke *et al.* 1993; Katz *et al.* 1995). Suicidal ideation was scored as positive if participants endorsed any of the following three items: 'felt life was not worth living', 'recurring thoughts of suicide' and 'wishing yourself were dead'. Physical health status was based on self-report using the RAND-36 Health Status Inventory (Adams & Benson, 1990; Hays, 1998), which has good reliability (Boyle *et al.* 1995; Feeny *et al.* 2002; Thoma *et al.* 2005) and validity (Feeny, 2002; Thoma *et al.* 2005). This measure specifically asked about current or past problems with

physical health (hypertension and asthma). The prevalence percentage rates for current/life-time hypertension and asthma were 5.5/10.1 and 7.3/14.1 respectively.

Potential confounders and missing values

Confounders included mother's age, parity, income, education and marital status, poverty level at delivery and child's sex, race, age at last interview and poverty level at age 7. Poverty level (Hardy *et al.* 1998) represents the ratio of the mother's annualized income to the poverty level based on the *Annual Statistical Supplement to the Social Security Bulletin* (US DHHS, 1993). All confounders, except the age of the child, were based on the mother's self-report. The child's age was calculated from birth dates. The frequency of missing data in this sample was negligible (less than 0.1%) for all control variables except for the mother's income and education attainment (2.2% missing).

Data analysis

First, each dichotomous perinatal factor was examined in separate maximum likelihood logistic regressions predicting each of the psychological and physical health outcomes. Second, to evaluate the magnitude of the association between co-morbid psychological and physical health problems, odds ratios (ORs) were obtained, using logistic regression without covariates and then with adjustment for potential confounders. Third, four dichotomized perinatal risk factors that the offspring had were examined as to whether they would predict their future co-morbidity of psychological and physical health problems status. Polytomous logistic regression analyses were conducted, using the Statistical Analysis System Categorical Model (CATMOD; SAS Institute, Cary, NC, USA), with four groups according to their co-morbid status: one with neither psychological nor physical health problems, one with physical health problems only, one with psychological health problems only, and one with co-morbidity. Finally, to explore whether different etiological processes exist during pregnancy, cluster analysis was used to examine whether these four perinatal indicators revealed natural clusters within a data-set that would otherwise not be apparent. Empirical internal testing indicates that the procedure is fairly

robust to violations of the assumptions of independence and the normal distribution (Norusis, 2004), but when the skewness or kurtosis were more than 1.96, we normalized the distribution using a logarithmic transformation. Cluster analysis began with the construction of a cluster features (CF) tree, which provides a capsule summary of the data file. The leaf nodes of the CF tree were then grouped using an agglomerative clustering algorithm. To determine which number of clusters was 'best', each of these cluster solutions was compared using Schwarz's Bayesian Criterion (BIC) as the clustering criterion. When clusters were extracted, we first evaluated the mean (s.d.) of the four perinatal characteristics in each cluster and then examined the overall difference in the risk for health problems among the cluster groups. The pairwise comparisons of the rates of each problem were further examined.

RESULTS

Descriptions of demographics and the four perinatal problems

The subjects were predominately black (82%), and approximately half (53%) were female. During the initial interviews between 1960 and 1964, mothers reported average ages of 24.9 (s.d. = 7.1) years and three (s.d. = 2.3) children. During the delivery interview, 86.2% of mothers reported completing primary school education, 22% high school, and less than 1% college. At the last interview (1992–1994), the mean age (s.d.) was 29 (2.5) years. Family income (s.d.) was US\$33682 (US\$24675). More than 60% of offspring had completed high school.

With the exception of the Apgar scores, perinatal factors were moderately to highly correlated. Birthweight was correlated with gestational length, head circumference and Apgar scores at 0.46, 0.79 and 0.17 respectively. Gestational length was correlated with head circumference and Apgar scores at 0.44 and 0.10 respectively. Apgar scores and head circumference were correlated at 0.11. Furthermore, tetrachoric correlations for dichotomous indices of perinatal risks were evaluated. LBW was correlated with preterm birth, small head circumference and low Apgar scores at 0.66, 0.89 and 0.33 respectively. Gestational length was

Table 1. Rates (*n*) and crude and adjusted risk of psychological and physical health problems in offspring by four perinatal risk factors (*n* = 1525)

Perinatal problems	Adult health problems ^a							
	Depression		Suicidal ideation		Hypertension		Asthma	
	% (<i>n</i>)	OR (95% CI) aOR (95% CI)	% (<i>n</i>)	OR (95% CI) aOR (95% CI)	% (<i>n</i>)	OR (95% CI) aOR (95% CI)	% (<i>n</i>)	OR (95% CI) aOR (95% CI)
Apgar scores								
Low (81)	8.6 (7)	4.1 (1.7–9.4)****	8.5 (7)	2.1 (0.95–4.8)*	7.1 (6)	1.4 (0.59–3.3)	7.4 (6)	1.0 (0.33–2.1)
Normal (1444)	2.3 (33)	3.5 (1.3–9.6)***	4.2 (61)	1.9 (0.73–5.0)	5.2 (75)	1.0 (0.36–3.2)	7.3 (105)	0.8 (0.28–2.3)
Head circumference								
Small (151)	6.6 (10)	3.1 (1.5–6.8)****	8.0 (12)	2.1 (1.1–3.9)**	10.7 (16)	2.5 (1.4–4.3)****	9.3 (14)	1.4 (0.76–2.4)
Normal (1374)	2.2 (30)	3.9 (1.7–8.6)****	4.0 (55)	2.4 (1.1–4.6)**	4.9 (67)	2.4 (1.3–4.4)***	7.0 (96)	1.4 (0.75–2.6)
Birthweight								
Low (235)	6.0 (14)	3.1 (1.6–6.0)****	6.8 (16)	2.1 (1.2–3.7)***	8.5 (20)	1.8 (1.1–3.0)**	10.2 (24)	1.6 (1.0–2.6)**
Normal (1290)	2.0 (26)	3.3 (1.6–6.8)****	4.0 (51)	2.1 (1.2–3.9)***	5.0 (64)	1.7 (1.0–2.8)**	6.7 (86)	1.5 (0.87–2.4)
Birth term								
Preterm (261)	5.0 (13)	2.4 (1.2–4.7)***	7.2 (18)	2.0 (1.1–3.4)***	8.4 (22)	1.8 (1.1–2.9)**	9.1 (24)	1.2 (0.75–2.0)
Normal (1264)	2.1 (27)	2.6 (1.3–5.4)***	3.8 (49)	2.1 (1.2–3.9)***	4.9 (62)	1.7 (1.0–3.0)**	7.0 (88)	1.4 (0.82–2.3)

OR, Odds ratio; aOR, adjusted odds ratio (mother's age, education, parity, marital status and poverty level at delivery and child's sex, age at last interview, race at poverty level at age 7 were adjusted); CI, confidence interval.

Perinatal problem scales (Apgar scores, preterm birth, birthweight, and head circumference) were created according to the traditional cut-off points: Apgar scores (low < 7 and normal ≥ 7); preterm birth (preterm < 37 completed weeks and normal ≥ 37 completed weeks); birthweight (low < 2500 g and normal ≥ 2500 g); and head circumference (small < 32 cm and normal ≥ 32 cm).

^a The General Health Questionnaire-28 (Goldberg, 1978) was used for depression and suicidal ideation and the RAND-36 Health Status Inventory (Adams & Benson, 1990) was used for current history of asthma and hypertension, with 1 representing the presence and 0 the absence of the illness.

* 0.10 < *p* < 0.05, ** 0.05 < *p* < 0.01, *** 0.01 < *p* < 0.001, **** 0.001 < *p* < 0.0001.

correlated with head circumference and Apgar scores at 0.70 and 0.24 respectively. Apgar scores and head circumference were correlated at 0.39.

Risk of psychological and physical health problems by perinatal problems

The relationships between the different perinatal problems singly and adult health problems are shown in Table 1. Offspring with small head circumference had an almost fourfold increased risk for depression (*p* = 0.003) and more than a twofold increased risk for suicidal ideation (*p* = 0.02) and hypertension (*p* = 0.01). Those with a low Apgar score had a more than threefold increased risk for depression (*p* = 0.01). Those with LBW had a threefold increased risk for depression (*p* < 0.001) and a twofold increased risk for suicidal ideation (*p* = 0.012) and hypertension (*p* = 0.08). Preterm birth was associated with an approximately twofold increased risk for depression (*p* = 0.01), suicidal ideation (*p* = 0.01) and hypertension (*p* = 0.05). None of the four risk factors elevated the risk for asthma after controlling for potential

confounders. Nearly all relationships remained equivalent or grew stronger when confounders were controlled for.†

Risk of co-morbid psychological and physical health problems

Table 2 shows the magnitude of co-morbid psychological and physical health problems after controlling for confounders. Offspring with depression had an elevated risk for co-morbid hypertension [OR 7.1, 95% confidence interval (CI) 3.2–15.8] and asthma (OR 3.3, 95% CI 1.4–8.0). Those with suicidal ideation also had an elevated risk for co-morbid hypertension (OR 2.7, 95% CI 1.3–6.2) and asthma (OR 2.5, 95% CI 1.2–5.3).

† Depression symptomatology was also evaluated based on a continuous measure of the GHQ depression subscale among offspring with and without a perinatal problem. Those with a problem relative to those without showed significantly higher depression scores after controlling for potential confounders. Adjusted means for the depression score for those with and without a perinatal problem and its associated *p* value were as follows (small head: 8.72 v. 8.07, *F* = 5.50, *p* = 0.007; low Apgar scores: 8.85 v. 8.09, *F* = 5.42, *p* = 0.02; LBW: 8.67 v. 8.03, *F* = 5.89, *p* = 0.001; and preterm birth: 8.58 v. 8.04, *F* = 5.56, *p* = 0.006).

Table 2. Risk of co-occurring psychological problems among those with hypertension and asthma in young adulthood ($n = 1525$)

	Psychological health problems					
	Depression			Suicidal ideation		
	No ^a ($n = 1485$) %	Yes ^b ($n = 40$) %	OR (95% CI) aOR (95% CI)	No ^a ($n = 1458$) %	Yes ^b ($n = 67$) %	OR (95% CI) aOR (95% CI)
Physical health problems						
Hypertension	4.9	27.5	7.4 (3.6–15.5)**** 7.1 (3.2–15.8)****	5.0	14.9	3.3 (1.6–6.8)*** 2.7 (1.3–6.2)*
Asthma	7.0	17.5	2.8 (1.2–6.5)** 3.3 (1.4–8.0)**	7.0	13.4	2.1 (1.0–4.3)* 2.8 (1.4–5.8)**

OR, Odds ratio; CI, confidence interval; aOR, adjusted odds ratio (mother's age, education, parity, marital status and poverty level at delivery and child's sex, age at last interview, race at poverty level at age 7 were adjusted).

^a 'No' for depression denotes the absence of a clinically significant condition, and 'No' for suicidal ideation denotes the absence of suicidal thoughts and ideas.

^b 'Yes' for depression denotes a clinically significant condition, and 'Yes' for suicidal ideation denotes the presence of suicidal thoughts and ideas.

* $0.05 \geq p > 0.01$, ** $0.01 \geq p > 0.001$, *** $0.001 \geq p > 0.0001$, **** $0.0001 \geq p$.

Table 3 shows whether perinatal risk factors might be associated with an increased risk for future health problems (psychological health problems only, physical health problems only, and co-morbidity). The offspring with LBW were at sevenfold, those with small head circumference at fivefold, and those with preterm birth at nearly fourfold increased risk for being in the co-morbid depression and hypertension group than in neither health problem group. However, only small head circumference was associated with a marginal increased risk for co-morbid hypertension and suicidal ideation (OR 4.4, $p = 0.08$). There was an approximately two- to threefold increased risk for offspring with a perinatal risk factor being in the psychological health problem (depression and suicidal ideation) only group. There was no evidence that perinatal risk factors increase the risk for co-morbid psychological problems and asthma.

Clusters of perinatal conditions in our sample and the risk for future psychological and physical health problems, and co-morbidity

Prior to running the cluster analysis, we tested whether each continuous perinatal indicator was normally distributed. Except for the Apgar scores, skewness and kurtosis were less than 1.96, meeting requirement for normal distribution. Apgar scores were normalized as they were negatively skewed (-2.84 , $s.d. = 0.063$) and had a high kurtosis (11.2 , $s.d. = 0.13$).

Cluster analysis identified four clusters. Cluster 1 consisted of offspring with preterm LBW, represented by low birthweight (mean 1869 g, range 1794–1945), preterm birth (mean 32.3 weeks, range 31.6–33.0), small head size (mean 29.9 cm, range 29.5–30.2) and a lower Apgar score (mean 7.67, range 7.34–8.0). Cluster 2 appears to consist of offspring with small gestational age (SGA), represented by lower birthweight (mean 2799 g, range 2774–2823 g), smaller head circumference (mean 32.8 cm, range 32.7–32.9) for gestational age (mean 38.1 weeks, range 37.9–38.3), and a normal Apgar score (mean 9.1, range 9.1–9.2). Cluster 3 appears to consist of offspring with fetal distress alone represented by normal birthweight (mean 3116 g, range 3025–3208), normal gestational age (mean 40.5 weeks, range 39.9–41.0), normal head circumference (mean 34.2 cm, range 33.9–34.5), but a very low Apgar score (mean 5.32, range 5.93–5.71). Finally, Cluster 4 appears to consist of offspring with optimal perinatal outcomes, normal birthweight (mean 3424 g, range 3399–3449), full-term gestational age (mean 40.4 weeks, range 40.3–40.6), normal head circumference (mean 34.7 cm, range 34.6–34.8) and a normal Apgar score (mean 9.13, range 9.09–9.18).

Table 4 shows the rates of depression, suicidal ideation, hypertension and asthma and the overall and pairwise differences in these rates among the cluster groups. Depression, suicidal

Table 3. Rates (n) and the risk for co-morbid physical (hypertension and asthma) and psychological (depression and suicidal ideation) health problems by perinatal problems

	% (n)	aOR (95% CI)	% (n)	aOR (95% CI)	% (n)	aOR (95% CI)	% (n)	aOR (95% CI)
(A) Co-morbid psychological problems and hypertension								
	Neither depression nor hypertension (n=1412)		Hypertension only (n=72)		Depression only (n=29)		Co-morbid (n=11)	
Head circumference	9.1 (128)	reference	18.1 (13)	1.5 (0.3-8.8)	20.7 (6)	2.5 (0.5-12.5)	36.4 (4)	5.6 (1.2-26.3)**
Low Apgar scores	5.0 (70)	reference	5.6 (4)	0.7 (0.1-7.7)	17.2 (5)	3.0 (0.3-34.5)	18.2 (2)	2.6 (0.3-23.1)
LBW	14.7 (207)	reference	19.4 (14)	1.3 (0.7-2.5)	27.6 (8)	2.7 (1.1-6.4)**	54.5 (6)	5.4 (1.5-20.0)***
Preterm birth	16.4 (232)	reference	22.2 (16)	1.3 (0.7-2.6)	27.6 (8)	2.4 (1.0-5.7)**	45.5 (5)	3.6 (0.92-14.1)*
	Neither suicidal ideation nor hypertension (n=1384)		Hypertension only (n=73)		Suicide ideation only (n=57)		Co-morbid (n=10)	
Head circumference	8.9 (123)	reference	19.2 (14)	2.6 (1.3-14.9)**	19.3 (11)	3.1 (1.5-6.2)***	30.0 (3)	4.4 (0.8-5.5)*
Low Apgar scores	4.9 (68)	reference	6.8 (5)	1.2 (0.4-4.4)	12.3 (7)	2.9 (1.2-7.3)**	10.0 (1)	N.E.
LBW	14.5 (200)	reference	23.3 (17)	1.8 (1.0-3.2)*	26.3 (15)	2.3 (1.2-4.5)***	30.0 (3)	1.9 (0.7-7.2)
Preterm birth	16.1 (223)	reference	26.0 (19)	1.9 (1.1-3.4)	29.8 (17)	2.5 (1.4-4.9)***	20.0 (2)	0.6 (0.04-5.0)
(B) Co-morbid psychological problems and asthma								
	Neither depression nor asthma (n=1381)		Asthma only (n=104)		Depression only (n=33)		Co-morbid (n=7)	
Head circumference	9.4 (129)	reference	11.9 (13)	1.3 (0.8-1.4)	25.0 (8)	4.5 (1.9-11.1)****	14.3 (1)	2.2 (0.1-50.0)
Low Apgar scores	5.0 (69)	reference	4.8 (5)	0.8 (0.3-2.4)	21.2 (7)	4.5 (1.6-12.5)***	0 (0)	N.E.
LBW	14.5 (200)	reference	20.2 (21)	1.3 (0.8-2.3)	33.3 (11)	3.2 (1.5-7.1)***	42.9 (3)	4.0 (0.8-20.0)
Preterm birth	16.5 (228)	reference	19.2 (20)	1.3 (0.8-2.2)	33.3 (11)	2.9 (1.3-6.3)**	28.6 (2)	2.5 (0.05-14.3)
	Neither suicidal ideation nor asthma (n=1357)		Asthma only (n=101)		Suicide ideation only (n=57)		Co-morbid (n=10)	
Head circumference	9.2 (125)	reference	11.9 (12)	1.3 (0.7-2.6)	21.1 (12)	3.1 (1.5-6.3)****	20.0 (2)	3.9 (0.5-12.5)
Low Apgar scores	5.1 (69)	reference	4.0 (4)	0.7 (0.9-2.6)	12.3 (7)	2.3 (0.9-6.3)***	10.0 (1)	2.9 (0.3-20.0)
LBW	14.4 (195)	reference	21.8 (22)	1.5 (0.9-2.6)	28.1 (16)	2.4 (1.3-4.6)***	20.0 (3)	1.3 (0.3-6.3)
Preterm birth	16.4 (222)	reference	19.8 (20)	1.4 (0.8-2.4)	29.8 (17)	2.3 (1.2-4.4)***	20.0 (2)	1.4 (0.3-7.1)

aOR, Adjusted odds ratio; CI, confidence interval; LBW, low birthweight; N.E., not estimable.

Apgar scores (low <7 and normal ≥7); preterm birth (pre-term <37 completed weeks and normal ≥37 completed weeks); birthweight (low <2500 g and normal ≥2500 g); and head circumference (small <32 cm and normal ≥32 cm). Polytomous logistic regression with potential confounders was used for statistical analysis.

* 0.10 > p > 0.05, ** 0.05 ≥ p > 0.01, *** 0.01 ≥ p > 0.001, **** 0.001 ≥ p > 0.0001.

ideation and asthma showed a significant overall group difference, with the preterm LBW group having the highest rate (7.8, 9.7 and 12.6% respectively), but the overall difference was only marginally significant for asthma ($p=0.08$) and hypertension ($p=0.09$) after adjusting for confounders. The last row shows the results from the pairwise comparisons. The preterm LBW group constantly showed poorer health outcomes. Moreover, the offspring with preterm LBW had a substantial increased risk for being in the co-morbid depression and hypertension group (OR 6.3, 95% CI 1.2-24.0), the co-morbid depression and asthma group (OR 10.2, 95% CI 1.8-58.7), the co-morbid suicidal ideation and hypertension group (OR 2.1, 95% CI 1.0-4.5), and the co-morbid

suicidal ideation and asthma group (OR 2.0, 95% CI 1.0-3.8) than being in neither health problem group (table available upon request).

DISCUSSION

Our data are consistent with and expand results from prior studies, and provide four main findings. First, perinatal problems, namely LBW, preterm birth and small head circumference, were associated with a significantly increased risk for both psychological and physical health problems in adulthood. Second, there was a significantly increased risk of co-occurring depression and physical problems in adult offspring. Third, single perinatal risk factors, namely LBW, preterm birth and small head

Table 4. Rates of depression, suicidal ideation, hypertension and asthma among the four cluster groups and overall unadjusted and adjusted difference in rates of these problems among the four cluster groups

	Depression (<i>n</i> =691) % (<i>n</i>)	Suicidal ideation (<i>n</i> =650) % (<i>n</i>)	Hypertension (<i>n</i> =81) % (<i>n</i>)	Asthma (<i>n</i> =103) % (<i>n</i>)
Clusters ^a				
1. Normal	2.3 (16)	3.8 (26)	4.8 (33)	7.4 (50)
2. SGA	1.8 (12)	4.2 (27)	5.8 (38)	7.2 (47)
3. Fetal distress	4.9 (4)	4.9 (4)	2.5 (2)	1.2 (1)
4. Preterm LBW	7.8 (8)	9.7 (10)	10.7 (11)	12.6 (13)
Unadjusted overall difference, Wald χ^2	14.1, <i>p</i> =0.003	7.9, <i>p</i> =0.052	7.6, <i>p</i> =0.056	8.8, <i>p</i> =0.033
Adjusted ^b overall difference, Wald χ^2	11.0, <i>p</i> =0.01	7.8, <i>p</i> =0.053	6.2, <i>p</i> =0.09	6.8, <i>p</i> =0.08
Pairwise difference ^c	4 > 1, 2	4 > 1	4 > 1, 2, 3	4 > 1, 2 > 3

SGA, Small for gestation age; LBW, low birthweight.

^a The four cluster groups were based on cluster analysis by four perinatal risk factors (birthweight, gestational age at delivery, 5-min Apgar scores, and head circumference at birth). The overall difference among the four groups was based on logistic regression.

^b Adjusted for mother's age, education, parity, marital status and poverty level at delivery and child's sex, age at the last interview, race, and poverty level at age 7.

^c The number represents the cluster group. Only significant (*p*<0.05) difference between the two cluster groups on the rate of each health problem is listed.

circumference, appear to be associated with an increased risk for co-morbid psychological problems and hypertension, but not co-morbid psychological problems and asthma. Fourth, there was initial evidence for differential etiological processes for preterm LBW, SGA and fetal distress, with preterm LBW having substantial adverse influence over adult health and co-morbidity.

We have demonstrated that perinatal problems have long-term effects on both psychological and physical health problems in adulthood, even after controlling for a wide range of confounders. Although the fetal origin hypothesis (Barker, 1995) suggested a link between malnutrition during pregnancy and later physical health conditions, especially heart disease (Leon *et al.* 1998; Eriksson *et al.* 2001; Barker & Lackland, 2003), only recently have studies begun to examine the possibility that perinatal problems influence the development of adult psychopathology, such as depression (Thompson *et al.* 2001, Gale & Martyn, 2004; Wiles *et al.* 2005; Nomura *et al.* in press) and schizophrenia (Hulshoff *et al.* 2000). However, to our knowledge no studies have examined psychological and physical health problems simultaneously. Our findings extend prior studies by showing that linkages between perinatal problems and adult illness are not limited to physical or psychological health problems.

The level of co-morbidity we found between psychological and physical health problems was striking. If offspring had depression, the risk of co-morbid hypertension increased more than sevenfold and asthma increased threefold. Similarly, if offspring had suicidal ideation, the risk for co-morbid hypertension and asthma increased more than twofold. Both psychological and physical health conditions were based on current problems. We therefore do not know if one problem proceeded to the other. It should be noted, however, that the rates of current hypertension (5.5%) and asthma (7.3%) were low in this relatively young adult population (mean age 29). As rates of depression (2.6%) and suicidal ideation (4.4%) were also low, the number of offspring with co-morbid conditions was small. Thus, the results need to be interpreted with great caution. The risk for co-morbid depression and hypertension by perinatal problems was evident but we found no evidence for co-morbid psychological problems and asthma by perinatal problems. Because significant co-morbidity between asthma and depression (OR 3.3, *p*=0.007) and between asthma and suicidal ideation (OR 2.8, *p*=0.005) were documented, this indicates that there may be different pathogenesis for the co-morbid asthma and psychological problems. Postnatal environment is one strong possibility for the increased risk. However, investigating this is

beyond the scope of the current study. Future studies could consider adverse postnatal psychosocial environment such as exposure to smoking and poverty (Rimington *et al.* 2001; Bender, 2006) as potential factors for the co-morbid psychosocial problems and asthma. We should remind readers that because low Apgar scores were only associated with an increased risk for psychological but not for physical health problems, the rate of low Apgar scores among the co-morbid group was very low. This prevented us from estimating the risk for co-morbid problems by low Apgar scores. However, it appears that Apgar scores, among the four perinatal measures that we examined, may influence adult health outcomes differently. This is consistent with the result based on cluster analysis; one cluster group that comprised offspring with optimal perinatal indicators except for low Apgar scores (the fetal distress group) had low rates of hypertension (1%) and asthma (2.5%) but relatively high rates of depression (5%) and suicidal ideation (5%). Tetrachoric correlation based on dichotomous Apgar scores (low *v.* normal) and any other perinatal risk indices were relatively low. This also suggests that a low Apgar score might have different etiological implications for the future illness, and including information on Apgar scores in an examination of fetal origin of adult illness and separating different processes might prove fruitful in future studies.

Note also that female offspring were more likely to be followed up through their adulthood ($p < 0.001$). It is well known that the risk for depression is generally higher among females and the risk for hypertension is higher among males. It is possible that our sample had an artificially higher rate of depression and lower rate of hypertension. Although there were no gender differences in rates of depression (2.8% for females *v.* 2.4% for males, $p = 0.69$) and hypertension (5.9% for females *v.* 5.2% for males, $p = 0.54$) in our sample, we do not know how the differential attrition by gender may have influenced our results, especially in the co-morbidity analysis.

There is considerable speculation about the mechanism for CNS alterations following fetal exposure to stress during the critical period of development. Some studies have uncovered evidence of brain injury among those with

preterm birth (Lemons *et al.* 1981; Garcia-Alix *et al.* 2004). Although preliminary, we attempted to elucidate different etiological processes based on the four perinatal scores (birthweight, gestational age, head circumference, and Apgar scores). As stated earlier, those with fetal distress, characterized by significantly low Apgar scores but otherwise normal perinatal scores, for example, had few problems with physical health but relatively high rates of psychological problems, while SGA appeared to have few psychological problems but moderately high rates of asthma (6%) and hypertension (7%). One cluster group (referred to as preterm LBW) that consisted of offspring with preterm birth and LBW together with small head circumference and low Apgar scores appeared to have the worst outcomes of all. We attempted to see whether this condition discriminated offspring with increased risk for co-morbid psychological and physical health problems and found a substantially increased risk for co-morbid depression and hypertension (OR 6.3, $p = 0.02$), co-morbid depression and asthma (OR 10.2, $p = 0.01$), co-morbid suicidal ideation and hypertension (OR 2.1, $p = 0.045$), and co-morbid suicidal ideation and asthma (OR 2.0, $p = 0.045$). Despite this significant co-morbidity between psychological problems and asthma, perinatal problems singly did not predict an increased risk for asthma and, consequently, there was no evidence for co-morbidity between psychological problems and asthma due to perinatal problems. Examining the underlying processes and mechanisms rather than single risk factors may prove more relevant to refining our understanding of perinatal health. One drawback in this approach, however, is that setting up the criteria we found based on cluster analysis could be arbitrary and therefore hinder systematic replication. Nevertheless, to our knowledge this is the first study to combine perinatal information and attempt to delineate underlying problems to evaluate the risk for co-morbid psychological and physical health problems. The influence of perinatal problems singly on increased risk for adult health status demonstrated that perinatal problems in general have a lingering consequence on poor health even in adulthood, whereas cluster analysis, delineating possible underlying etiological processes, helped us to understand who are at

highest risk for co-morbid psychological and physical health problems in adulthood. Future studies should consider examining the actual biological mechanism during pregnancy that leads the offspring to have an adverse condition such as preterm LBW.

Our study has several methodological strengths. The second-generation cohort was prospectively and systematically followed from birth and studied longitudinally for almost 30 years. The subjects were randomly selected regional (inner-city Baltimore, USA) representatives who sought prenatal care at Johns Hopkins Hospital rather than a clinical sample with either serious psychopathology or medical illness. Thus, co-morbid conditions reported here had less chance of being artificially inflated and have higher external validity. Three of the four perinatal problems (birthweight, Apgar scores, and head circumference) were recorded by a research nurse at the time of delivery rather than based on mother's retrospective report.

This study also has limitations. First, as stated earlier, we had no direct measure of HPA-axis functioning, although we speculate that this is the mechanism through which perinatal problems increase the risk for co-morbid conditions. However, our preterm LBW group, extracted by cluster analysis, could be viewed as the one with the most adverse influence on the HPA axis. We still recognize that this approach does not directly reveal HPA-axis function. Second, both psychological and physical health problems in adulthood were based on self-report checklist inventories, not clinical diagnostic assessments. Although the validity of the instruments is widely supported, there were no external measures to validate the reliability of the self-report. Self-report on physical health conditions often leads to under-reporting. For example, we considered diabetes as another physical health problem, but this was not possible because of its low prevalence ($n=14$, 0.9%). In the general population, the prevalence in this age group ranges from 3.0% to 4.0% (Connolly, 2006) and even two to six times higher among the disadvantaged (Liao *et al.* 2004; Mainous *et al.* 2004; Mensah *et al.* 2005; Russo *et al.* 2006), which describes our sample. Moreover, it is possible that individuals with psychological problems may have systematically

over-reported asthma and hypertension, while those without these problems may have under-reported them. This would artificially inflate/deflate the increased risk for co-morbid psychological and physical health problems. Third, there is some evidence that parental depression increases the risk for offspring's affective and anxiety disorders and cardiovascular illness (Weissman *et al.* 2006; Nomura *et al.* in press). Although this may or may not be through a different pathway, it is an important factor to consider when we examine the risk for co-morbid psychological and physical problems. Unfortunately, our data do not include the mother's psychological status and therefore we were not able to control for this effect. Fourth, the number of offspring with co-morbid psychological and physical health problems ranged from 7 to 11 (Table 3). Although the magnitude of associations was significant, the small cell sizes contributed to wide confidence intervals. Thus, the results should be interpreted with caution. Future studies using samples with greater rates of physical health problems would allow us to re-examine the risk for co-morbidity between psychological and physical conditions by perinatal problems. These findings warrant replication and extension to include investigation of the perinatal precursors to compromised psychological and physical health across the life cycle.

ACKNOWLEDGMENTS

This study was supported by R03MH067761 (Y.N.) from the NIMH and the Young Investigator Award (Y.N.) from the National Alliance for Research on Schizophrenia and Depression (NARSAD). We thank Janet Hardy and Sam Shapiro, principal investigators of the Pathways to Adulthood Study, for allowing us to use their data and we are grateful to the mothers and their children who participated in the study. We also thank Priya J. Wickramaratne for her constructive advice on an early version of the manuscript. Finally, we thank Claude M. Chemtob and Jeffery H. Newcorn for their support of this project.

DECLARATION OF INTEREST

None.

REFERENCES

- Adams, P. F. & Benson, V. (1990). Current estimates from the National Health Interview Survey, 1989. *Vital and Health Statistics*, Series 10, No. 176. DHHS Publication No. (PHS) 90-1504. United States Government Printing Office: Washington, DC.
- Allen, N. B., Lewinshon, P. M. & Seeley, J. R. (1998). Prenatal and perinatal influences on risk for psychopathology in childhood and adolescence. *Development and Psychopathology* **10**, 513–529.
- Anda, R. F., Williamson, D. F., Jones, D., Maccera, C., Eaker, E., Glassman, A. H. & Marks, J. (1993). Depressed affect, hopelessness, and the risk for ischemic heart disease in a cohort of US adults. *Epidemiology* **4**, 285–294.
- Barker, D. J. P. (1995). Fetal origins of coronary heart disease. *British Medical Journal* **311**, 171–174.
- Barker, D. J. P., Eriksson, J. G., Forsen, T. & Osmond, C. (2002). Fetal origins of adult disease: strength of effects and biological basis. *International Journal of Epidemiology* **31**, 1235–1239.
- Barker, D. J. P. & Lackland, D. T. (2003). Prenatal influences on stroke mortality in England and Wales. *Stroke* **34**, 1598–1603.
- Bender, B. G. (2006). Risk taking, depression, adherence, and symptom control in adolescents and young adult with asthma. *American Journal of Respiratory and Critical Care Medicine* **173**, 953–957.
- Boyle, M. H., Furlong, W., Feeny, D., Torrance, G. W. & Hatcher, J. (1995). Reliability of the Health Utilities Index – Mark III used in the 1991 cycle 6 General Social Survey Health Questionnaire. *Quality of Life Research* **4**, 249–257.
- Broadhead, W. E., Blazer, D. G., George, L. K. & Tse, C. K. (1990). Depression, disability days, and days lost from work in a prospective epidemiologic survey. *Journal of the American Medical Association* **264**, 2524–2538.
- Chuckley, S. (1996). The neuroendocrinology of depression. *British Medical Bulletin* **52**, 597–617.
- Clarke, D. M., Smith, G. C. & Herrman, H. (1993). A comparative study of screening instruments for mental disorders in general hospital patients. *International Journal of Psychiatry and Medicine* **23**, 323–337.
- Connolly, V. (2006). Diabetes, deprivation and outcomes in the wealthy world. *Diabetes Voice* **51**, 39–40.
- Cox, D., Lewis, G., Stuart, G. & Murphy, K. (2002). A cross-sectional study of the prevalence of psychopathology in adults with congenital heart disease. *Journal of Psychosomatic Research* **52**, 65–68.
- Davies, A. A., Davey-Smith, G., May, M. T. & Ben-Shlomo, Y. (2006). Association between birth weight and blood pressure is robust, amplifies with age, and may be underestimated. *Hypertension* **48**, 431–436.
- Druss, B. & Pincus, H. (2000). Suicidal ideation and suicide attempts in general medical illness. *Archives of Internal Medicine* **160**, 1522–1526.
- Eriksson, J. G., Forsen, T., Tuomilehto, J., Osmond, C. & Barker, D. J. P. (2001). Early growth and coronary heart disease in later life: longitudinal study. *British Medical Journal* **22**, 949–953.
- Fan, A. P. & Eaton, W. W. (2001). Longitudinal study assessing the joint effects of socio-economic status and birth risks on adult emotional and nervous conditions. *British Journal of Psychiatry. Supplement* **40**, s78–s83.
- Feeny, D. H., Furlong, W. J., Torrance, G. W. & Barr R. D. (2002). Multiattribute and single-attribute utility functions for the Health Utilities Index Mark 3 system. *Medical Care* **40**, 113–228.
- Feeny, D. (2002). Commentary on Jack Dowie, ‘Decision validity should determine whether a generic or condition-specific HRQOL measure is used in health care decisions.’ *Health Economics* **11**, 13–16.
- Fulop, G., Strain, J. J., Fahs, M. C., Hammer, J. S. & Lyons, J. S. (1988). Medical disorders associated with psychiatric comorbidity and prolonged hospital stay. *Hospital and Community Psychiatry* **40**, 80–82.
- Gale, C. R. & Martyn, C. N. (2004). Birth weight and later risk of depression in a national birth cohort. *British Journal of Psychiatry* **184**, 28–33.
- Garcia-Alix, A., Saenz-de Pipaon, M., Martine, M., Salas-Hernandez, S. & Quero, J. (2004). Ability of neonatal head circumference to predict long-term neurodevelopmental outcome. *Revista de Neurologia* **39**, 548–554.
- Gillaspay, S. R., Hoff, A. L., Mullins, L. L., van Pelt, J. C. & Chaney, J. M. (2002). Psychological distress in high-risk youth with asthma. *Journal of Pediatrics and Psychology* **27**, 363–371.
- Glassman, A. H., Bigger, J. T., Gaffney, M., Shapiro, P. A. & Swenson, J. R. (2006). Onset of major depression associated with acute coronary syndromes: relationship of onset, major depressive disorder history, and episode severity to sertraline benefit. *Archives of General Psychiatry* **63**, 283–288.
- Goldberg, D. (1978). *Manual of the General Health Questionnaire*. NFER: Windsor.
- Goldberg, D. P. (1972). *The Detection of Psychiatric Illness by Questionnaire: A Technique for the Identification and Assessment of Non-psychotic Psychiatric Illness*. Oxford University Press: London.
- Goodwin, R. D., Fergusson, D. M. & Horwood, L. J. (2004). Asthma and depressive and anxiety disorders: results of a 21-year longitudinal study. *Psychological Medicine* **34**, 1465–1474.
- Hales, C. N., Barker, D. J., Clark, P. M., Cox, L. J., Fall, C., Osmond, C. & Winter, P. D. (1991). Fetal and infant growth and impaired glucose tolerance at age 64. *British Medical Journal* **303**, 1019–1022.
- Hardy, J. B. (2003). The Collaborative Perinatal Project: lessons and legacy. *Annals of Epidemiology* **13**, 303–311.
- Hardy, J. B., Astone, N. M., Brooks-Gunn, J., Shapiro, S. & Miller, T. L. (1998). Like mother, like child: intergenerational patterns of age at first birth and associations with childhood and adolescent characteristics and adult outcomes in the second generation. *Developmental Psychology* **34**, 1269–1273.
- Hardy, J. B., Shapiro, S., Mellits, D., Skinner, E. A., Astone, N. M., Ensminger, M., Laveist, T., Baumgardner, R. A. & Starfield, B. H. (1997). Self-sufficiency at age 27 to 33 years: factors present between birth and 18 years that predict educational attainment among children born to inner-city families. *Pediatrics* **99**, 80–87.
- Hays, R. D. (1998). *Rand Health Status Inventory*. The Psychological Corporation/Harcourt Brace & Company: Orlando.
- Heim, C. & Nemeroff, C. B. (2001). The role of childhood trauma in the neurobiology of mood and anxiety disorders: preclinical and clinical studies. *Biological Psychiatry* **49**, 1023–1039.
- Hulshoff Pol, H. E., Hoek, H. W., Susser, E., Brown, A. S., Dingemans, A., Schnack, H. G., van Haren, N. E., Pereira Ramos, L. M., Gispens-de Wied, C. C. & Kahn, R. S. (2000). Prenatal exposure to famine and brain morphology in schizophrenia. *American Journal of Psychiatry* **157**, 1170–1172.
- Hurwitz, E. L. & Morgenstern, H. (1999). Cross-sectional associations of asthma, hay fever, and other allergies with major depression and low-back pain among adults aged 20–39 years in the United States. *American Journal of Epidemiology* **150**, 1107–1116.
- Huxley, R., Neil, A. & Collins, R. (2002). Unraveling the fetal origins hypothesis: is there really an inverse association between birthweight and subsequent blood pressure? *Lancet* **360**, 659–665.
- Jones, C. A., Feeny, D. & Eng, K. (2005). Test–retest reliability of Health Utilities Index scores: evidence from hip fracture. *International Journal of Technology Assessment in Health Care* **21**, 393–398.
- Katz, R., Stephen, J., Shaw, B. F., Matthew, A., Newman, F. & Rosenbluth, M. (1995). The East York Health Needs Study: prevalence of DSM-III-R psychiatric disorder in a sample of Canadian women. *British Journal of Psychiatry* **166**, 100–106.
- Kramer, R. A., Warner, V., Olfson, M., Banks, C. M., Chaput, F. & Weissman, M. M. (1998). General medical problems among the offspring of depressed parents: a 10-year follow-up. *Journal of the American Academy of Child and Adolescent Psychiatry* **37**, 1–9.
- Lee, S., Yu, H., Wing, Y., Chen, C., Lee, A. M., Lee, D. T. S., Chen, C., Lin, K. & Weiss, M. G. (2000). Psychiatric morbidity and

- illness experience of primary care patients with chronic fatigue in Hong Kong. *American Journal of Psychiatry* **157**, 380–384.
- Lemons, J. A., Schreiner, R. I. & Grasham, E. L.** (1981). Relationship of brain weight to head circumference in early infancy. *Human Biology* **53**, 351–354.
- Leon, D. A., Litchell, H. O. & Vegero, D.** (1998). Reduced fetal growth rate and increased risk of death from ischaemic heart disease: cohort study of 15000 Swedish men and women born 1915–1929. *British Medical Journal* **317**, 241–245.
- Liao, Y., Tucker, P., Okoro, C. A., Giles, W. H., Mokdad, A. H. & Harris, V. B.** (2004). REACH 2010 surveillance for health status in minority communities: United States, 2001–2002. *Morbidity and Mortality Weekly Report. Surveillance Summaries* **53**, 1–36.
- Mainous, A. G., King, D. E., Garr, D. R. & Pearson, W. S.** (2004). Race, rural residence, and control of diabetes and hypertension. *Annals of Family Medicine* **2**, 563–568.
- Meaney, M. J.** (2001). Maternal care, gene expression, and the transmission of individual differences in stress reactivity across generations. *Annual Review of Neuroscience* **24**, 1161–1192.
- Mensah, G. A., Mokdad, A. H., Ford, E. S., Greenlund, K. J. & Croft, J. B.** (2005). State of disparities in cardiovascular health in the United States. *Epidemiology* **111**, 1233–1241.
- Musselman, D. L., Evans, D. L. & Nemeroff, C. B.** (1998). The relationship of depression to cardiovascular disease. *Archives of General Psychiatry* **55**, 580–592.
- Nichols, P. L. & Chen, T. C.** (1981). *Minimal Brain Dysfunction: A Prospective Study*. Lawrence Erlbaum Associates: Hillsdale, NJ.
- Nomura, Y., Wickramaratne, P. J., Pilowsky, D. J., Bruder, B., Newcorn, J. H., Fifer, W. P., Brooks-Gunn, J. & Weissman, M. M.** (in press). Low birth weight and risk of affective disorders and selected medical illness in offspring at high and low risk for depression. *Comprehensive Psychiatry*.
- Norusis, M.** (2004). *SPSS 13.0 Statistical Procedures Companion*. Prentice Hall, Inc.: Upper Saddle-River, NJ.
- Ortega, A. N., McQuaid, E. L., Canino, G., Goodwin, R. D. & Fritz, G. K.** (2004). Comorbidity of asthma and anxiety and depression in Puerto Rican children. *Psychosomatics* **45**, 93–99.
- Patton, G. C., Coffey, C., Carlin, J. B., Olsson, C. A. & Morley, R.** (2004). Prematurity at birth and adolescent depressive disorder. *British Journal of Psychiatry* **184**, 446–447.
- Rimington, L. D., Davies, D. H., Lowe, D., Pearson, M. G., Russo, C. A., Andrew, R. M. & Coffey, R. M.** (2001). Relationship between anxiety, depression and morbidity in adult asthma patients. *Thorax* **56**, 266–271.
- Russo, C. A., Andrews, R. M. & Coffey, R. M.** (2006). Racial and ethnic disparities in potentially preventable hospitalizations, 2003. *Healthcare Cost and Utilization Project* **10**, 1–9.
- Sanchez, M. M., Ladd, C. O. & Plotsky, P. M.** (2001). Early adverse experience as developmental risk factor for late psychopathology: evidence from rodent and primate models. *Developmental Psychopathology* **13**, 419–449.
- Sandman, C. A., Wadhwa, P. D., Chicz-DeMet, A., Porto, M. & Garite, T. J.** (1998). Maternal corticotropin-releasing hormone and habituation in the human fetus. *Developmental Psychobiology* **34**, 163–173.
- Thoma, A., Sprague, S., Veltr, K., Duku, E. & Furlong, W.** (2005). Methodology and measurement properties of health-related quality of life instruments: a prospective study of patients undergoing breast reduction surgery. *Health and Quality of Life Outcomes* **3**, 44.
- Thompson, C., Syddall, H., Rodin, I., Osmond, C. & Barker, D. J.** (2001). Birth weight and the risk of depressive disorder in later life. *British Journal of Psychiatry* **179**, 450–455.
- US DHHS** (1993). *Social Security Bulletin. Annual Statistical Supplement, 1993*, pp. 346–348. US Department of Health and Human Services: Washington, DC.
- Vaag, A., Jensen, C. B., Poulsen, P., Brons, C., Pilgaard, K., Grunnet, L., Vielwerth, S. & Alibegovic, A.** (2006). Metabolic aspects of insulin resistance in individuals born small for gestational age. *Hormone Research* **65**, s137–s143.
- van Melle, J. P., de Jonge, P., Spijkerman, T. A., Tijssen, J. G., Ormel, J., van Veldhuisen, D. J., van den Brink, R. H. & van den Berg, M. P.** (2004). Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis. *Psychosomatic Medicine* **66**, 814–822.
- Veening, M. A., Van Weissenbruch, M. M. & Delemarre-Van De Waal, H. A.** (2002). Glucose tolerance, insulin sensitivity, and insulin secretion in children born small for gestational age. *Journal of Clinical Endocrinology and Metabolism* **87**, 4657–4661.
- Vila, G., Nollet-Clemencon, C., de Blic, J., Mouren-Simeoni, M. C. & Scheinmann, P.** (2000). Prevalence of DSM IV anxiety and affective disorders in a pediatric population of asthmatic children and adolescents. *Journal of Affective Disorders* **58**, 223–231.
- Wadhwa, P. D., Porto, M., Garite, T. J., Chicz-DeMet, A. & Sandman, C. A.** (1998). Maternal corticotropin-releasing hormone levels in the early third trimester predict length of gestation in human pregnancy. *American Journal of Obstetrics and Gynecology* **179**, 1079–1085.
- Wadhwa, P. D., Sandman, C. A. & Garite, T. J.** (2001). The neurobiology of stress in human pregnancy: implications for prematurity and development of the fetal central nervous system. *Progress in Brain Research* **133**, 131–142.
- Wamboldt, M. Z., Hewitt, J. K., Schmitz, S., Wamboldt, F. S., Rasanen, M., Koskenvuo, M., Romanav, K., Varjonen, J. & Kaprio, J.** (2000). Familial association between allergic disorders and depression in adult Finnish twins. *American Journal of Medical Genetics (Neuropsychiatric Genetics)* **96**, 146–153.
- Ward, A. J.** (1991). Prenatal stress and childhood psychopathology. *Child Psychiatry and Human Development* **22**, 97–110.
- Weissman, M. M., Wickramaratne, P. J., Nomura, Y., Warner, V., Pilowsky, D. J. & Vedeli, H.** (2006). Offspring of depressed parents: 20 years later. *American Journal of Psychiatry* **163**, 1001–1008.
- Wells, K. B., Golding, J. M. & Burnam, M. A.** (1988). Psychiatric disorder in a sample of the general population with and without chronic medical conditions. *American Journal of Psychiatry* **145**, 976–981.
- Wiles, N. J., Peters, T. J., Leon, D. A. & Lewis G.** (2005). Birth weight and psychological distress at age 45–51 years. *British Journal of Psychiatry* **187**, 21–28.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.